


Rathen 21.09.09

# Top Quark Physics at ATLAS

## from Cross-Section to New Physics

Valentina Ferrara




# Curriculum Vitae

- ☒ Laurea Triennale
  - “*Studies of the Read Out System of the ATLAS Data Acquisition System*”
  - Fernanda Pastore - Roma Tre
  - Livio Mapelli - CERN


- ☒ Laurea Magistrale - major subject: particle physics
  - “*The ATLAS discovery potential for a charged slepton as next-to-lightest supersymmetric particle*”
  - Domizia Orestano - Roma Tre
  - Klaus Moenig - DESY

- ☐ Top Physics with the ATLAS detector at LHC
  - ...



# Contents




- Standard Model predictions for Top Physics at LHC
  - Top Physics phenomenology
- Measurements with first data: top-pair production cross-section
  - ATLAS trigger efficiency from data
- A handle on new physics
- Summary





# Standard Model Predictions



## Production @ LHC



gluon-gluon fusion 90%

quark-antiquark annihilation 10%

## Cross-Section



- $\sigma_{tt} (\sqrt{s} = 14 \text{ TeV}) \approx 886 \text{ pb}$
- $\sigma_{tt} (\sqrt{s} = 10 \text{ TeV}) \approx 403 \text{ pb}$
- $\sigma_{tt} (\sqrt{s} = 7 \text{ TeV}) \approx 161 \text{ pb}$

This translates into ( $\sqrt{s} = 14 \text{ TeV}$ ):


- ~ $88.6 \times 10^3$  top pairs in  $100 \text{ pb}^{-1}$  of data
- ~ $10^7$  top pairs per year before selection

A precision era will begin for top quark physics at the LHC!


(\*) arXiv:0907.2527

# Top Quark Phenomenology

The top quark decays rapidly without forming hadrons, and almost exclusively through the single mode  $t \rightarrow Wb$



Name	Signature	BR	xsec at 10 TeV
Fully Hadronic	jets	45.7%	191.5 pb
Lepton + Jets	$e + \text{jets}$	17.2%	71.9 pb
	$\mu + \text{jets}$	17.2%	71.9 pb
Dilepton	$e\mu + \text{jets}$	3.18%	13.3 pb
	$\mu\mu + \text{jets}$	1.59%	6.67 pb
	$ee + \text{jets}$	1.59%	6.67 pb
Tau + Jets	$\tau + \text{jets}$	9.49%	39.8 pb
Lepton + Tau	$\tau + e/\mu + \text{jets}$	3.54%	14.8 pb
Tau + Tau	$\tau + \tau + \text{jets}$	0.49%	2.06 pb
total	all	100%	419 pb



Semi-leptonic decay mode

Signature:

- one lepton
- missing energy from the neutrino
- two jets from the W boson
- two jets from the b quark

# Determination of the Cross-Section

Given the high statistics which will be available even in the initial phase of the LHC, the top-pair cross section measurement can be performed **relatively fast** and even **with an imperfectly calibrated** detector

## Counting method:

number of observed events  
meeting the selection criteria  
of top-event signature

$$\sigma = \frac{N_{sig}}{L \times \epsilon} = \frac{N_{obs} - N_{bkg}}{L \times \epsilon}$$


geometrical acceptance  
trigger efficiency  
event selection efficiency

number of background  
events estimated from  
Monte Carlo simulation  
and/or data samples


■ Trigger efficiencies determined from simulation  
represent a dangerous source of systematic uncertainty and  
should be determined directly from data

→ **Tag&Probe technique**


# Trigger efficiency from Data



i) first consider a sample of events that pass the three ATLAS trigger levels with at least a muon with transverse momentum higher than 20 GeV




ii) for every muon that has fired the trigger (“**Tag**” muon) we require a second muon (“**Probe**”) to be identified offline together with some identification conditions on the Z particle (e.g. the invariant mass of the pair must match the mass of the Z)



iii) The trigger efficiency is defined by the frequency with which the **Probe** muon in this sample also passed the trigger selection



# Trigger efficiency from Data - Simulations



Z-boson decays and top-pair decays have different features...  
can we apply efficiencies measured on  $Z\mu\mu$  events to top-pair events?



# Trigger efficiency from Data - Simulations



Parametrizing the efficiency with more than one variable allows to apply a weight on top-pair events according to the muon **kinematic values** and **position** in the detector


# A Handle on New Physics

Are the top pairs produced solely by gluon-gluon fusion and quark-antiquark annihilation?  
Is there some **heavy object X** which decays into top quarks, thus enhancing the observed rate over QCD predictions?

Is the cross section in line with predictions regardless of the final state or is some **non-standard decay** of top somehow modifying the mix?

Do the other properties of the top quark give preference to the Standard Model top quark hypothesis over more exotic scenarios?

D0 and CDF have already initiated some of these studies but are limited by the statistics acquired



# Summary

- The precise measurement of the top quark properties is an important task for the LHC
- Cross Section measurements are an important test of possible new production mechanisms (e.g. decay of heavier objects such as the top quark super-partner or Kaluza-Klein resonances)
- New physics may also modify the cross section times branching ratio differently in various decay channels (e.g. supersymmetric channels with charged Higgs  $t \rightarrow H^+ b$  or with super-partner of the top quark  $t \rightarrow t \chi^0$ )